

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188	
<p>The public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden, to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.</p>					
1. REPORT DATE 28 August 2017		2. REPORT TYPE Briefing Charts		3. DATES COVERED (From - To) 01 August 2017 - 31 August 2017	
4. TITLE AND SUBTITLE Variable Weight Fractional Collisions for Multiple Species Mixtures				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) Robert Martin				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER Q1AM	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Air Force Research Laboratory (AFMC) AFRL/RQRS 1 Ara Drive Edwards AFB, CA 93524-7013				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) Air Force Research Laboratory (AFMC) AFRL/RQR 5 Pollux Drive Edwards AFB, CA 93524-7048				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S) AFRL-RQ-ED-VG-2017-205	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for Public Release; Distribution Unlimited. PA Clearance Number: 17517 Clearance Date: 22 August 2017					
13. SUPPLEMENTARY NOTES For presentation at Direct Simulation Monte Carlo 2017; Santa Fe, NM, USA; 28 August 2017 The U.S. Government is joint author of the work and has the right to use, modify, reproduce, release, perform, display, or disclose the work.					
14. ABSTRACT Viewgraph/Briefing Charts					
15. SUBJECT TERMS N/A					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON
a. REPORT	b. ABSTRACT	c. THIS PAGE			Robert Martin
Unclassified	Unclassified	Unclassified	SAR	86	19b. TELEPHONE NUMBER (Include area code) N/A

VARIABLE WEIGHT FRACTIONAL COLLISIONS FOR MULTIPLE SPECIES MIXTURES

Robert Martin

IN-SPACE PROPULSION BRANCH,
AIR FORCE RESEARCH LABORATORY,
EDWARDS AIR FORCE BASE, CA USA

Direct Simulation Monte Carlo, 2017
Distribution Statement A: Approved for public release;
Distribution is Unlimited; PA #17517



U.S. AIR FORCE






- 1 BACKGROUND & REVIEW OF METHOD
- 2 MULTI-SPECIES TEST CASES
- 3 FUTURE WORK
- 4 CONCLUSION



Field-Reversed Configuration:

- Concept from Fusion Energy
 - Scaled Down for Propulsion

RP3X FRC Thruster






FRC THRUSTER OVERVIEW


Field-Reversed Configuration:

- Concept from Fusion Energy
 - Scaled Down for Propulsion
- Electrodeless
 - Limits Erosion
 - Enables Flexible Fuels



Pancotti, et al, "Adaptive Electric Propulsion for ISRU Missions", 20th Adv. Space Prop., 11/2014

RP3X FRC Thruster






Field-Reversed Configuration:

- Concept from Fusion Energy
 - Scaled Down for Propulsion
- Electrodeless
 - Limits Erosion
 - Enables Flexible Fuels
- Pulsed Operation
 - Tunable Thrust/ISP
 - Coupled Dynamics

RP3X FRC Thruster






Field-Reversed Configuration:

- Concept from Fusion Energy
 - Scaled Down for Propulsion
- Electrodeless
 - Limits Erosion
 - Enables Flexible Fuels
- Pulsed Operation
 - Tunable Thrust/ISP
 - Coupled Dynamics
- Challenges Common to Pulsed EM

RP3X FRC Thruster






Field-Reversed Configuration:

- Concept from Fusion Energy
 - Scaled Down for Propulsion
- Electrodeless
 - Limits Erosion
 - Enables Flexible Fuels
- Pulsed Operation
 - Tunable Thrust/ISP
 - Coupled Dynamics
- Challenges Common to Pulsed EM

Complex to Design

RP3X FRC Thruster






Field-Reversed Configuration:

- Concept from Fusion Energy
 - Scaled Down for Propulsion
- Electrodeless
 - Limits Erosion
 - Enables Flexible Fuels
- Pulsed Operation
 - Tunable Thrust/ISP
 - Coupled Dynamics
- Challenges Common to Pulsed EM

Complex to Design
Significant Modeling Challenge


RP3X FRC Thruster





PHASES OF FRC OPERATION

Dominant Physics Varies with Cycle:





Adapted from "Annular FRC" PhD Proposal, C. (Niemela) Hill



PHASES OF FRC OPERATION

Dominant Physics Varies with Cycle:

- 1 Neutral Fill
- Rarefied Kinetic Flow




Adapted from "Annular FRC" PhD Proposal, C. (Niemela) Hill



PHASES OF FRC OPERATION

Dominant Physics Varies with Cycle:

- ① Neutral Fill
 - Rarefied Kinetic Flow
- ② Preionization Chemistry
 - CR-Excitation/Ionization




Adapted from "Annular FRC" PhD Proposal, C. (Niemela) Hill



PHASES OF FRC OPERATION

Dominant Physics Varies with Cycle:

- ① Neutral Fill
 - Rarefied Kinetic Flow
- ② Preionization Chemistry
 - CR-Excitation/Ionization
- ③ Driver Pulse
 - Ionization+Electromagnetics




Adapted from "Annular FRC" PhD Proposal, C. (Niemela) Hill




PHASES OF FRC OPERATION

Dominant Physics Varies with Cycle:

- 1 Neutral Fill
 - Rarefied Kinetic Flow
- 2 Preionization Chemistry
 - CR-Excitation/Ionization
- 3 Driver Pulse
 - Ionization+Electromagnetics
- 4 Field Reversal
 - Magnetic Reconnection



<https://astrobear.pas.rochester.edu/trac/wiki/AstroBearProjects/resistiveMHD>





Adapted from "Annular FRC" PhD Proposal, C. (Niemela) Hill



PHASES OF FRC OPERATION

Dominant Physics Varies with Cycle:

- 1 Neutral Fill
 - Rarefied Kinetic Flow
- 2 Preionization Chemistry
 - CR-Excitation/Ionization
- 3 Driver Pulse
 - Ionization+Electromagnetics
- 4 Field Reversal
 - Magnetic Reconnection
- 5 Plasmoid Ejection
 - $\vec{j} \times \vec{B}$, Neutral Entrainment



Adapted from "Annular FRC" PhD Proposal, C. (Niemela) Hill




PHASES OF FRC OPERATION

Dominant Physics Varies with Cycle:

- ① Neutral Fill
 - Rarefied Kinetic Flow
- ② Preionization Chemistry
 - CR-Excitation/Ionization
- ③ Driver Pulse
 - Ionization+Electromagnetics
- ④ Field Reversal
 - Magnetic Reconnection
- ⑤ Plasmoid Ejection
 - $\vec{j} \times \vec{B}$, Neutral Entrainment

Continuous Cycle: ⑤ impacts ①



Adapted from "Annular FRC" PhD Proposal, C. (Niemela) Hill



IMPORTANCE OF COLLISION PHYSICS


Important Collisions in Spacecraft Propulsion:

- Discharge and Breakdown in FRC
- Collisional Radiative Cooling/Ionization
- Combustion Chemistry


Common Features in Spacecraft Collisions:

- Relevant Densities Spanning Many Orders of Magnitude — 6+
- Transitions from Collisional to Collisionless
- Tiny Early e^- or Radical Populations Critical to Induction Delay
- Many types of Inelastic Collisions with Unknown Effects on Distribution Shapes

Shock Ionization



Kapper & Cambier, J. Appl. Phys. 109, (2011)





IMPORTANCE OF COLLISION PHYSICS

Important Collisions in Spacecraft Propulsion:


- Discharge and Breakdown in FRC
- Collisional Radiative Cooling/Ionization
- Combustion Chemistry

Common Features in Spacecraft Collisions:


- Relevant Densities Spanning Many Orders of Magnitude — 6+
- Transitions from Collisional to Collisionless
- Tiny Early e^- or Radical Populations Critical to Induction Delay
- Many types of Inelastic Collisions with Unknown Effects on Distribution Shapes

Need Low Noise & High Dynamic Range
Collision Algorithms

Shock Ionization



Kapper & Cambier, J. Appl. Phys. 109, (2011)





Previous Collision Methods:


- Monte Carlo Collisions (MCC)
 - Particles Collide with Background “Fluid”
 - Often Used in Plasma/PIC Simulation
 - Ion- e^- Collisions Assume Stationary Ions
 - No Conservation/Detailed Balance
- Direct Simulation Monte Carlo Collisions (DSMC)
 - Most Modern Versions use No-Time Counter (NTC) Method
 - Conservative/Reversible Collision
 - Satisfies Detailed Balance
 - Subset of Possible Collisions Sampled
 - Random Selection vs Z_{ij} for All/Nothing Collision

All Random Flip vs Number of Collisions: $Z_{ij} = \frac{n_i n_j}{2} \langle \sigma v \rangle dt$



Continuum to Discrete Representation:

- Many Particles \rightsquigarrow Continuous Distribution





VARIABLE WEIGHTS FOR DYNAMIC RANGE

Continuum to Discrete Representation:

- Many Particles \rightsquigarrow Continuous Distribution
- Discretized VDF Yields Vlasov
But Collision Integral Still a Problem





Continuum to Discrete Representation:

- Many Particles \rightsquigarrow Continuous Distribution
- Discretized VDF Yields Vlasov
But Collision Integral Still a Problem
- Particle Methods VDF to Delta Function Set
- Collisions between Discrete Velocities






VARIABLE WEIGHTS FOR DYNAMIC RANGE

Continuum to Discrete Representation:

- Many Particles \rightsquigarrow Continuous Distribution
- Discretized VDF Yields Vlasov
But Collision Integral Still a Problem
- Particle Methods VDF to Delta Function Set
- Collisions between Discrete Velocities
- But Poorly Resolved Tail
(Tail Critical to Inelastic Collisions)





VARIABLE WEIGHTS FOR DYNAMIC RANGE

Continuum to Discrete Representation:

- Many Particles \leadsto Continuous Distribution
- Discretized VDF Yields Vlasov
But Collision Integral Still a Problem
- Particle Methods VDF to Delta Function Set
- Collisions between Discrete Velocities
- But Poorly Resolved Tail
(Tail Critical to Inelastic Collisions)
- Variable Weights Permit Extra DOF in Tails






Continuum to Discrete Representation:

- Many Particles \rightsquigarrow Continuous Distribution
- Discretized VDF Yields Vlasov
But Collision Integral Still a Problem
- Particle Methods VDF to Delta Function Set
- Collisions between Discrete Velocities
- But Poorly Resolved Tail
(Tail Critical to Inelastic Collisions)
- Variable Weights Permit Extra DOF in Tails

Variable Weight “All-or-Nothing” Collisions?






VARIABLE WEIGHTS FOR DYNAMIC RANGE

Continuum to Discrete Representation:

- Many Particles \rightsquigarrow Continuous Distribution
- Discretized VDF Yields Vlasov
But Collision Integral Still a Problem
- Particle Methods VDF to Delta Function Set
- Collisions between Discrete Velocities
- But Poorly Resolved Tail
(Tail Critical to Inelastic Collisions)
- Variable Weights Permit Extra DOF in Tails

Variable Weight “All-or-Nothing” Collisions?






Continuum to Discrete Representation:

- Many Particles \rightsquigarrow Continuous Distribution
- Discretized VDF Yields Vlasov
But Collision Integral Still a Problem
- Particle Methods VDF to Delta Function Set
- Collisions between Discrete Velocities
- But Poorly Resolved Tail
(Tail Critical to Inelastic Collisions)
- Variable Weights Permit Extra DOF in Tails

Variable Weight “All-or-Nothing” Collisions?

Physically Inconsistent!

(Mixing Violates Momentum/Energy Conservation)





Stochastic Weighted Particle Method:

- Developed by Rjasanow & Wagner

Attempted Collisions/Cell:

$$\nu = f(2\bar{w} - w_{min})N_p(N_p - 1) \langle \sigma v \rangle^{max} dt$$

Select Pair (i,j) if:

$$\text{Rand} < \frac{w_i + w_j - w_{min}}{N_p(N_p - 1)(2\bar{w} - w_{min})}$$

-or-

$$\text{Rand} < \frac{w_i + w_j - w_{min}}{(2w_{max} - w_{min})}$$

Collide If:

$$\text{Rand} < \frac{\langle \sigma v \rangle_{ij}}{\langle \sigma v \rangle^{max}} \frac{f \max(w_i, w_j)}{w_i + w_j - w_{min}}$$

Perform Standard VHS Collisions

Generate/Modify Particles with:

$$\pm \Delta w / f = \pm \min(w_i, w_j) / f$$

Update $\langle \sigma v \rangle^{max}$



Stochastic Weighted Particle Method:

- Developed by Rjasanow & Wagner
- Adapted as Modified NTC/MCF

Attempted Collisions/Cell:

$$\nu = f(2\bar{w} - w_{min})N_p(N_p - 1) \langle \sigma v \rangle^{max} dt$$

Select Pair (i,j) if:

$$Rand < \frac{w_i + w_j - w_{min}}{N_p(N_p - 1)(2\bar{w} - w_{min})}$$

-or-

$$Rand < \frac{w_i + w_j - w_{min}}{(2w_{max} - w_{min})}$$

Collide If:

$$Rand < \frac{\langle \sigma v \rangle_{ij}}{\langle \sigma v \rangle^{max}} \frac{f \max(w_i, w_j)}{w_i + w_j - w_{min}}$$

Perform Standard VHS Collisions

Generate/Modify Particles with:

$$\pm \Delta w / f = \pm \min(w_i, w_j) / f$$

Update $\langle \sigma v \rangle^{max}$



Stochastic Weighted Particle Method:

- Developed by Rjasanow & Wagner
- Adapted as Modified NTC/MCF
- Assures Post-Collision $w_i \geq 0$

Attempted Collisions/Cell:

$$\nu = f(2\bar{w} - w_{min})N_p(N_p - 1) \langle \sigma v \rangle^{max} dt$$

Select Pair (i,j) if:

$$\text{Rand} < \frac{w_i + w_j - w_{min}}{N_p(N_p - 1)(2\bar{w} - w_{min})}$$

-or-

$$\text{Rand} < \frac{w_i + w_j - w_{min}}{(2w_{max} - w_{min})}$$

Collide If:

$$\text{Rand} < \frac{\langle \sigma v \rangle_{ij}}{\langle \sigma v \rangle^{max}} \frac{f \max(w_i, w_j)}{w_i + w_j - w_{min}}$$

Perform Standard VHS Collisions

Generate/Modify Particles with:

$$\pm \Delta w / f = \pm \min(w_i, w_j) / f$$

Update $\langle \sigma v \rangle^{max}$



Stochastic Weighted Particle Method:

- Developed by Rjasanow & Wagner
- Adapted as Modified NTC/MCF
- Assures Post-Collision $w_i \geq 0$
- Converges to NTC for $w_i = \text{const}$

Attempted Collisions/Cell:

$$\nu = f(2\bar{w} - w_{\min})N_p(N_p - 1) \langle \sigma v \rangle^{\max} dt$$

Select Pair (i,j) if:

$$\text{Rand} < \frac{w_i + w_j - w_{\min}}{N_p(N_p - 1)(2\bar{w} - w_{\min})}$$

-or-

$$\text{Rand} < \frac{w_i + w_j - w_{\min}}{(2w_{\max} - w_{\min})}$$

Collide If:

$$\text{Rand} < \frac{\langle \sigma v \rangle_{ij}}{\langle \sigma v \rangle^{\max}} \frac{f \max(w_i, w_j)}{w_i + w_j - w_{\min}}$$

Perform Standard VHS Collisions

Generate/Modify Particles with:

$$\pm \Delta w / f = \pm \min(w_i, w_j) / f$$

Update $\langle \sigma v \rangle^{\max}$



Stochastic Weighted Particle Method:

- Developed by Rjasanow & Wagner
- Adapted as Modified NTC/MCF
- Assures Post-Collision $w_i \geq 0$
- Converges to NTC for $w_i = \text{const}$
- Only Adds 1-particle/collision for $\Delta w = \min(w_i, w_j)$

Attempted Collisions/Cell:

$$\nu = f(2\bar{w} - w_{\min})N_p(N_p - 1) \langle \sigma v \rangle^{\max} dt$$

Select Pair (i,j) if:

$$\text{Rand} < \frac{w_i + w_j - w_{\min}}{N_p(N_p - 1)(2\bar{w} - w_{\min})}$$

-or-

$$\text{Rand} < \frac{w_i + w_j - w_{\min}}{(2w_{\max} - w_{\min})}$$

Collide If:

$$\text{Rand} < \frac{\langle \sigma v \rangle_{ij}}{\langle \sigma v \rangle^{\max}} \frac{f \max(w_i, w_j)}{w_i + w_j - w_{\min}}$$

Perform Standard VHS Collisions

Generate/Modify Particles with:

$$\pm \Delta w / f = \pm \min(w_i, w_j) / f$$

Update $\langle \sigma v \rangle^{\max}$




Stochastic Weighted Particle Method:

- Developed by Rjasanow & Wagner
- Adapted as Modified NTC/MCF
- Assures Post-Collision $w_i \geq 0$
- Converges to NTC for $w_i = \text{const}$
- Only Adds 1-particle/collision for $\Delta w = \min(w_i, w_j)$
- Adds 2-particles/collision for $\Delta w = \min(w_i, w_j)/f$

Attempted Collisions/Cell:

$$\nu = f(2\bar{w} - w_{\min})N_p(N_p - 1) \langle \sigma v \rangle^{\max} dt$$



Generate/Modify Particles with:

$$\pm \Delta w / f = \pm \min(w_i, w_j) / f$$

Update $\langle \sigma v \rangle^{\max}$




Stochastic Weighted Particle Method:

- Developed by Rjasanow & Wagner
- Adapted as Modified NTC/MCF
- Assures Post-Collision $w_i \geq 0$
- Converges to NTC for $w_i = \text{const}$
- Only Adds 1-particle/collision for $\Delta w = \min(w_i, w_j)$
- Adds 2-particles/collision for $\Delta w = \min(w_i, w_j)/f$
- **Still Requires Merge $w_i \neq \text{const}$**

Attempted Collisions/Cell:

$$\nu = f(2\bar{w} - w_{\min})N_p(N_p - 1) \langle \sigma v \rangle^{\max} dt$$



Generate/Modify Particles with:

$$\pm \Delta w / f = \pm \min(w_i, w_j) / f$$


Update $\langle \sigma v \rangle^{\max}$



REVIEW OF CONSERVATIVE MERGE

Merge to Pair → DOF for Conservation:

- $(n+2):2$ yields Exact Mass, Momentum, and Kinetic Energy Conservation
- Applied Spatially also Shown to Conserve Electrostatic Energy
- Though Energy Conserving, Still Thermalizes VDF



$$w_{cell} = \sum_i^{(n+2)} w_i$$

$$\vec{v} = \frac{1}{w_{cell}} \sum_i^{(n+2)} w_i \vec{v}_i$$

$$\overline{V^2} = \frac{1}{w_{cell}} \sum_i^{(n+2)} w_i \left(\vec{v}_i - \vec{v} \right)^2$$

$$w_{(a/b)} = w_m/2$$

$$\vec{v}_{(a/b)} = \vec{v} \pm \hat{\mathcal{R}} \sqrt{\overline{V^2}}$$

$$\text{Similarly: } \vec{x}_{(a/b)} = \vec{x} \pm \hat{\mathcal{R}} \sqrt{\overline{X^2}}$$




REVIEW OF CONSERVATIVE MERGE

Merge to Pair \rightarrow DOF for Conservation:

- $(n+2):2$ yields Exact Mass, Momentum, and Kinetic Energy Conservation
- Applied Spatially also Shown to Conserve Electrostatic Energy
- Though Energy Conserving, Still Thermalizes VDF

Selection of Near Neighbors in VDF Limits Thermalization



$$w_{cell} = \sum_i^{(n+2)} w_i$$

$$\vec{v} = \frac{1}{w_{cell}} \sum_i^{(n+2)} w_i \vec{v}_i$$

$$\overline{V^2} = \frac{1}{w_{cell}} \sum_i^{(n+2)} w_i \left(\vec{v}_i - \vec{v} \right)^2$$

$$w_{(a/b)} = w_m / 2$$

$$\vec{v}_{(a/b)} = \vec{v} \pm \hat{\mathcal{R}} \sqrt{\overline{V^2}}$$

$$\text{Similarly: } \vec{x}_{(a/b)} = \vec{x} \pm \hat{\mathcal{R}} \sqrt{\overline{X^2}}$$



REVIEW OF CONSERVATIVE MERGE


Merge to Pair → DOF for Conservation:

- $(n+2):2$ yields Exact Mass, Momentum, and Kinetic Energy Conservation
- Applied Spatially also Shown to Conserve Electrostatic Energy
- Though Energy Conserving, Still Thermalizes VDF

Selection of Near Neighbors in VDF Limits Thermalization

Merge via Separate Octree/Species
Only Change for Mixtures!

Octree Velocity Bins




Efficient Neighbor Selection



FROM RGD30: MACH 2 ARGON SHOCK

1D Normal Argon Shock Test

- Simple Verification vs. DS1V
- Initial Conditions:
 $T_0 = 293\text{K}$, $n_0 = 1\text{E}22/\text{m}^3$, $v_0 = 637.4(\text{m/s})$
- Initial Jump to Post-Shock at 1cm
- VHS Collisions:
 $T_{ref}=273\text{K}$, $d_{ref}=4.17\text{\AA}$, $\omega_{VHS}=0.81$






FROM RGD30: MACH 2 ARGON SHOCK

1D Normal Argon Shock Test

- Simple Verification vs. DS1V
- Initial Conditions:
 $T_0 = 293\text{K}$, $n_0 = 1\text{E}22/\text{m}^3$, $v_0 = 637.4(\text{m/s})$
- Initial Jump to Post-Shock at 1cm
- VHS Collisions:
 $T_{ref}=273\text{K}$, $d_{ref}=4.17\text{\AA}$, $\omega_{VHS}=0.81$
- Time Average:
 \bar{n} from $t \in [80, 100)\mu\text{s}$

TURF - SWPM+Octree



Target N/Cell Quadrupled per Line





FROM RGD30: MACH 2 ARGON SHOCK

1D Normal Argon Shock Test

- Simple Verification vs. DS1V
- Initial Conditions:
 $T_0 = 293\text{K}$, $n_0 = 1\text{E}22/\text{m}^3$, $v_0 = 637.4(\text{m/s})$
- Initial Jump to Post-Shock at 1cm
- VHS Collisions:
 $T_{ref}=273\text{K}$, $d_{ref}=4.17\text{\AA}$, $\omega_{VHS}=0.81$
- Time Average:
 \bar{n} from $t \in [80, 100)\mu\text{s}$
- Error (Normalized L_1):
 $err = |n - \bar{n}|/\bar{n}$
- Error Controlled: $err \propto \sqrt{N/cell}$

TURF - SWPM+Octree



Target N/Cell Quadrupled per Line




FROM RGD30: MACH 8 ARGON BOW SHOCK

2D Argon Shock Test

- Initial Conditions like $M=2$ Except:
 $v_0 = 2550\text{m/s}$
- Specular: $x=5 - 5.04\text{mm}$ with $y=\pm 2\text{mm}$
- Half Domain Modeled:
 $80\mu\text{m} \times 80\mu\text{m}$ Cells

TURF: n - Standard DSMC




TURF: n - SWPM+Octree



FROM RGD30: MACH 8 ARGON BOW SHOCK

2D Argon Shock Test

- Initial Conditions like $M=2$ Except:
 $v_0 = 2550\text{m/s}$
- Specular: $x=5 - 5.04\text{mm}$ with $y=\pm 2\text{mm}$
- Half Domain Modeled:
 $80\mu\text{m} \times 80\mu\text{m}$ Cells
- Time Average:
 \bar{n} from $t \in [80, 100)\mu\text{s}$
- SWPM Similar to Standard DSMC






FROM RGD30: MACH 8 ARGON BOW SHOCK


2D Argon Shock Test

- Initial Conditions like $M=2$ Except:
 $v_0 = 2550\text{m/s}$
- Specular: $x=5 - 5.04\text{mm}$ with $y=\pm 2\text{mm}$
- Half Domain Modeled:
 $80\mu\text{m} \times 80\mu\text{m}$ Cells
- Time Average:
 \bar{n} from $t \in [80, 100)\mu\text{s}$
- SWPM Similar to Standard DSMC
- Despite Different Np/Cell

TURF: n - Standard DSMC



TURF: n - SWPM+Octree
TURF Np/Cell - Standard DSMC




TURF Np/Cell - SWPM+Octree



HE:XE MIXTURE SHOCK

1D Normal He:Xe Shock Test

- Mach 3.89 with He:Xe of 97:3
(i.e. Bird '94 Fig 12.35)




Converged 100x Baseline



HE:XE MIXTURE SHOCK

1D Normal He:Xe Shock Test

- Mach 3.89 with He:Xe of 97:3 (i.e. Bird '94 Fig 12.35)
- Highlights Species Separation




Converged 100x Baseline



HE:XE MIXTURE SHOCK

1D Normal He:Xe Shock Test

- Mach 3.89 with He:Xe of 97:3 (i.e. Bird '94 Fig 12.35)
- Highlights Species Separation
- Separation Peak at $\rho_{He} \approx \rho_{Xe}$




Converged 100x Baseline



HE:Xe MIXTURE SHOCK

1D Normal He:Xe Shock Test

- Mach 3.89 with He:Xe of 97:3 (i.e. Bird '94 Fig 12.35)
- Highlights Species Separation
- Separation Peak at $\rho_{He} \approx \rho_{Xe}$
- DSMC needs 33x He:Xe Macroparticles (From m_{Xe}/m_{He})
- Reduced Particle Count Introduces Error




Comparison 10x vs. 100x



HE:Xe MIXTURE SHOCK

1D Normal He:Xe Shock Test

- Mach 3.89 with He:Xe of 97:3 (i.e. Bird '94 Fig 12.35)
- Highlights Species Separation
- Separation Peak at $\rho_{He} \approx \rho_{Xe}$
- DSMC needs 33x He:Xe Macroparticles (From m_{Xe}/m_{He})
- Reduced Particle Count Introduces Error
- Error in Time Average at Baseline




Comparison Baseline vs. 100x



HE:Xe MIXTURE SHOCK

1D Normal He:Xe Shock Test

- Mach 3.89 with He:Xe of 97:3 (i.e. Bird '94 Fig 12.35)
- Highlights Species Separation
- Separation Peak at $\rho_{He} \approx \rho_{Xe}$
- DSMC needs 33x He:Xe Macroparticles (From m_{Xe}/m_{He})
- Reduced Particle Count Introduces Error
- Error in Time Average at Baseline
- Instantaneous: Dramatic Noise in Xe



Instantaneous Baseline vs. 100x




HE:Xe MIXTURE SHOCK

1D Normal He:Xe Shock Test

- Mach 3.89 with He:Xe of 97:3 (i.e. Bird '94 Fig 12.35)
- Highlights Species Separation
- Separation Peak at $\rho_{He} \approx \rho_{Xe}$
- DSMC needs 33x He:Xe Macroparticles (From m_{Xe}/m_{He})
- Reduced Particle Count Introduces Error
- Error in Time Average at Baseline
- Instantaneous: Dramatic Noise in Xe

Noise Reduction via Variable Weights?




Instantaneous Baseline vs. 100x



He:Xe MIXTURE SHOCK - FRACTIONAL DSMC

1D He:Xe Shock with SWPM+Octrees

- Xe Noise Controlled by 1:1 Target
- He:Xe Noise Comparable




100x vs. FDSMC 1:1



He:Xe MIXTURE SHOCK - FRACTIONAL DSMC

1D He:Xe Shock with SWPM+Octrees

- Xe Noise Controlled by 1:1 Target
- He:Xe Noise Comparable
- Direct Noise Control by Target Ratio




100x vs. FDSMC 2:1



1D He:Xe Shock with SWPM+Octrees

- Xe Noise Controlled by 1:1 Target
- He:Xe Noise Comparable
- Direct Noise Control by Target Ratio




100x vs. FDSMC 4:1



1D He:Xe Shock with SWPM+Octrees

- Xe Noise Controlled by 1:1 Target
- He:Xe Noise Comparable
- Direct Noise Control by Target Ratio




100x vs. FDSMC 8:1



1D He:Xe Shock with SWPM+Octrees

- Xe Noise Controlled by 1:1 Target
- He:Xe Noise Comparable
- Direct Noise Control by Target Ratio




100x vs. FDSMC 16:1



HE:Xe MIXTURE SHOCK - FRACTIONAL DSMC

1D He:Xe Shock with SWPM+Octrees

- Xe Noise Controlled by 1:1 Target
- He:Xe Noise Comparable
- Direct Noise Control by Target Ratio
- Converged Error Finite! (1:1 Target)
- Error Source Still Unidentified...




Averaged 100x vs. FDSMC 1:1



1D He:Xe Shock with SWPM+Octrees

- Xe Noise Controlled by 1:1 Target
- He:Xe Noise Comparable
- Direct Noise Control by Target Ratio
- Converged Error Finite! (1:1 Target)
- Error Source Still Unidentified...
- Potentially Sensitivity to He Tails?
 - Merge Impacts Higher Moments
 - TBD Error vs. He-Noise Level
 - Improvement Merge to Preserve Tails
- Adaptation of SWMP Incorrect?
 - # of Collisions Sampled as WDF Varies?
 - Collision Pair Rule Wrong, $w_i \ll w_j$?




Averaged 100x vs. FDSMC 1:1



1D He:Xe Shock with SWPM+Octrees

- Xe Noise Controlled by 1:1 Target
- He:Xe Noise Comparable
- Direct Noise Control by Target Ratio
- Converged Error Finite! (1:1 Target)
- Error Source Still Unidentified...
- Potentially Sensitivity to He Tails?
 - Merge Impacts Higher Moments
 - TBD Error vs. He-Noise Level
 - Improvement Merge to Preserve Tails
- Adaptation of SWMP Incorrect?
 - # of Collisions Sampled as WDF Varies?
 - Collision Pair Rule Wrong, $w_i \ll w_j$?

Error Identification in Future Work




Averaged 100x vs. FDSMC 1:1



ROTHE'S HE:AR FREE-JET EXPERIMENT

- Helium-Argon Mixture
- Expanded through Nozzle to Vacuum
- e -Beam Concentration Measurements

Jet Flow Layout



$Re=533$





FIG. 11. Mole fraction of argon throughout the flow field of a free jet.




Case Tested in PhD & RGD27:

Continuum Boundary Conditions




Surrogate for FRC Injection



Dynamic Range \gg Shocks

Solution from DS2V



Solution from SPARTAN
(Navier-Stokes + Diffusion Velocity)



1 Wing Increases near Nozzle Edge






FIG. 11. Mole fraction of argon throughout the flow field of a free jet.

Experimental



DS2V




SPARTAN




FEATURES FOUND VIA SIMULATIONS


1 Wing Increases near Nozzle Edge



Experimental



DS2V




SPARTAN




FEATURES FOUND VIA SIMULATIONS


- 1 Wing Increases near Nozzle Edge
- 2 Lower Radial Boundary Edge Concentration



Experimental



DS2V




SPARTAN




FEATURES FOUND VIA SIMULATIONS




- 1 Wing Increases near Nozzle Edge
- 2 Lower Radial Boundary Edge Concentration
- 3 Deeper Jet Edge Concentration Drop



Experimental



DS2V



SPARTAN



- TURF Naturally 3D Cartesian
- Considered 2D-Axisymmetric TURF
- Simple DSMC but Merge Complex
(Conservation on $v_{\parallel} v_{\perp}$ -Quadrees?)


Standard 3D DSMC



CHALLENGES FOR JET IN TURF

- TURF Naturally 3D Cartesian
- Considered 2D-Axisymmetric TURF
- Simple DSMC but Merge Complex (Conservation on $v_{\parallel} v_{\perp}$ -Quadtrees?)
- Opted to Run Coarse Full 3D (Simplified Boundary Conditions)
- 3D Expensive at Tractable Resolution
- Added Collision Sub-Cells to TURF

Standard 3D DSMC






CHALLENGES FOR JET IN TURF

- TURF Naturally 3D Cartesian
- Considered 2D-Axisymmetric TURF
- Simple DSMC but Merge Complex (Conservation on $v_{\parallel} v_{\perp}$ -Quadtrees?)
- Opted to Run Coarse Full 3D (Simplified Boundary Conditions)
- 3D Expensive at Tractable Resolution
- Added Collision Sub-Cells to TURF
- Fractional DSMC Controls Np/Cell

Standard 3D DSMC




Multi-Species Fractional DSMC



CHALLENGES FOR JET IN TURF

- TURF Naturally 3D Cartesian
- Considered 2D-Axisymmetric TURF
- Simple DSMC but Merge Complex (Conservation on $v_{\parallel} v_{\perp}$ -Quadtrees?)
- Opted to Run Coarse Full 3D (Simplified Boundary Conditions)
- 3D Expensive at Tractable Resolution
- Added Collision Sub-Cells to TURF
- Fractional DSMC Controls Np/Cell
- Linear Density Obscures Results




Multi-Species Fractional DSMC



CHALLENGES FOR JET IN TURF

- TURF Naturally 3D Cartesian
- Considered 2D-Axisymmetric TURF
- Simple DSMC but Merge Complex (Conservation on $v_{||}v_{\perp}$ -Quadrees?)
- Opted to Run Coarse Full 3D (Simplified Boundary Conditions)
- 3D Expensive at Tractable Resolution
- Added Collision Sub-Cells to TURF
- Fractional DSMC Controls Np/Cell
- Linear Density Obscures Results
- Issue Clearer with NpAr/Cell






CHALLENGES FOR JET IN TURF

- TURF Naturally 3D Cartesian
- Considered 2D-Axisymmetric TURF
- Simple DSMC but Merge Complex (Conservation on $v_{\parallel} v_{\perp}$ -Quadrees?)
- Opted to Run Coarse Full 3D (Simplified Boundary Conditions)
- 3D Expensive at Tractable Resolution
- Added Collision Sub-Cells to TURF
- Fractional DSMC Controls Np/Cell
- Linear Density Obscures Results
- Issue Clearer with NpAr/Cell


X_{Ar} to RX-Plane \rightarrow for Detailed Results...






PRELIMINARY TURF RESULTS: HE:AR JET


1 Standard DSMC Poor Results



Experimental



DS2V




TURF

1:1 - Collision:Sample




PRELIMINARY TURF RESULTS: HE:AR JET


- 1 Standard DSMC Poor Results
- 2 2x2x2 Collision Cell Improves Standard DSMC



Experimental



DS2V




TURF

2x2x2:1 - Collision:Sample




PRELIMINARY TURF RESULTS: HE:AR JET


- 1 Standard DSMC Poor Results
- 2 2x2x2 Collision Cell Improves Standard DSMC
- 3 SWPM+Octree Significantly Better (2x2x2 Collision Cell)



Experimental



DS2V



TURF

SWPM+Octree (2x2x2 Collision Cells)



FUTURE DIRECTIONS: HYBRID ∂f

The ∂f -Boltzmann for $f = f^{eq} + f^{dev}$

- ∂f - Concept Old
- How to make ∂f cheaper than full- f
(Must adapt DOF Usage..?)




FUTURE DIRECTIONS: HYBRID ∂f

The ∂f -Boltzmann for $f = f^{eq} + f^{dev}$

- ∂f - Concept Old
- How to make ∂f cheaper than full- f
(Must adapt DOF Usage..?)
- Recent Progress using $\pm\delta$ -weight Particles

Strong Landau Damping



P-P:	$v+, w+$	\rightarrow	$v'+, w'+$
P-N:	$v+, w-$	\rightarrow	$2v+, v'-, w'-$
N-N:	$v-, w-$	\rightarrow	$2v-, 2w-, v'+, w'+$
P-M:	$M, v+$	\rightarrow	$M, w-, v'+, w'+$
N-M:	$M, v-$	\rightarrow	$M, w+, v'-, w'-$
M-M:		\rightarrow	\emptyset

Yan, JCP 309 (2016) 18-36




FUTURE DIRECTIONS: HYBRID ∂f

The ∂f -Boltzmann for $f = f^{eq} + f^{dev}$

- ∂f - Concept Old
- How to make ∂f cheaper than full- f (Must adapt DOF Usage..?)
- Recent Progress using $\pm\delta$ -weight Particles
- Requires Remapping due to Particle Growth

Strong Landau Damping



P-P:	$v+, w+$	\rightarrow	$v'+, w'+$
P-N:	$v+, w-$	\rightarrow	$2v+, v'-, w'-$
N-N:	$v-, w-$	\rightarrow	$2v-, 2w-, v'+, w'+$
P-M:	$M, v+$	\rightarrow	$M, w-, v'+, w'+$
N-M:	$M, v-$	\rightarrow	$M, w+, v'-, w'-$
M-M:		\rightarrow	\emptyset

Yan, JCP 309 (2016) 18-36




FUTURE DIRECTIONS: HYBRID ∂f

The ∂f -Boltzmann for $f = f^{eq} + f^{dev}$

- ∂f - Concept Old
- How to make ∂f cheaper than full- f (Must adapt DOF Usage..?)
- Recent Progress using $\pm\delta$ -weight Particles
- Requires Remapping due to Particle Growth
- SWPM has Similar Issue (Basis for Octree B2B Collisions)

Stochastic Weight Particle Method (SWPM)



$$w_i = w_i - \Delta w_{ij} \text{ \& } w_j = w_j - \Delta w_{ij}$$
$$w_{(N_p+1)} = \Delta w_{ij} \text{ \& } w_{(N_p+2)} = \Delta w_{ij}$$

+2 Particles/Collision

RGD30



FUTURE DIRECTIONS: HYBRID ∂f

The ∂f -Boltzmann for $f = f^{eq} + f^{dev}$

- ∂f - Concept Old
- How to make ∂f cheaper than full- f
(Must adapt DOF Usage..?)
- Recent Progress using $\pm\delta$ -weight Particles
- Requires Remapping due to Particle Growth
- SWPM has Similar Issue
(Basis for Octree B2B Collisions)
- Octree: Bin Moments, $M_i \rightarrow$ Particles

Octree ∂f -Boltzmann Bin-to-Bin Collisions






FUTURE DIRECTIONS: HYBRID ∂f

The ∂f -Boltzmann for $f = f^{eq} + f^{dev}$

- ∂f - Concept Old
- How to make ∂f cheaper than full- f (Must adapt DOF Usage..?)
- Recent Progress using $\pm\delta$ -weight Particles
- Requires Remapping due to Particle Growth
- SWPM has Similar Issue (Basis for Octree B2B Collisions)
- Octree: Bin Moments, $M_i \rightarrow$ Particles
- Root Bin Sum \rightarrow Equilibrium Moments

Octree ∂f -Boltzmann Bin-to-Bin Collisions






FUTURE DIRECTIONS: HYBRID ∂f

The ∂f -Boltzmann for $f = f^{eq} + f^{dev}$

- ∂f - Concept Old
- How to make ∂f cheaper than full- f (Must adapt DOF Usage..?)
- Recent Progress using $\pm\delta$ -weight Particles
- Requires Remapping due to Particle Growth
- SWPM has Similar Issue (Basis for Octree B2B Collisions)
- Octree: Bin Moments, $M_i \rightarrow$ Particles
- Root Bin Sum \rightarrow Equilibrium Moments
- ∂M_i from $M_i^{Bin} - M_i^{Eq}$

Octree ∂f -Boltzmann Bin-to-Bin Collisions






FUTURE DIRECTIONS: HYBRID ∂f

The ∂f -Boltzmann for $f = f^{eq} + f^{dev}$

- ∂f - Concept Old
- How to make ∂f cheaper than full- f (Must adapt DOF Usage..?)
- Recent Progress using $\pm\delta$ -weight Particles
- Requires Remapping due to Particle Growth
- SWPM has Similar Issue (Basis for Octree B2B Collisions)
- Octree: Bin Moments, $M_i \rightarrow$ Particles
- Root Bin Sum \rightarrow Equilibrium Moments
- ∂M_i from $M_i^{Bin} - M_i^{Eq}$
- B2B Collision Recast with δw Particles

Octree ∂f -Boltzmann Bin-to-Bin Collisions



Sample Collisions using
 Δw : for P-P, P-N, N-N, P-M, N-M
M-M: \emptyset




FUTURE DIRECTIONS: HYBRID ∂f

The ∂f -Boltzmann for $f = f^{eq} + f^{dev}$

- ∂f - Concept Old
- How to make ∂f cheaper than full- f
(Must adapt DOF Usage..?)
- Recent Progress using $\pm\delta$ -weight Particles
- Requires Remapping due to Particle Growth
- SWPM has Similar Issue
(Basis for Octree B2B Collisions)
- Octree: Bin Moments, $M_i \rightarrow$ Particles
- Root Bin Sum \rightarrow Equilibrium Moments
- ∂M_i from $M_i^{Bin} - M_i^{Eq}$
- B2B Collision Recast with δw Particles
- Collision Work $\propto \partial f$, not f

Octree ∂f -Boltzmann Bin-to-Bin Collisions



Sample Collisions using
 Δw : for P-P, P-N, N-N, P-M, N-M
 M-M: \emptyset




FUTURE DIRECTIONS: HYBRID ∂f

The ∂f -Boltzmann for $f = f^{eq} + f^{dev}$

- ∂f - Concept Old
- How to make ∂f cheaper than full- f (Must adapt DOF Usage..?)
- Recent Progress using $\pm\delta$ -weight Particles
- Requires Remapping due to Particle Growth
- SWPM has Similar Issue (Basis for Octree B2B Collisions)
- Octree: Bin Moments, $M_i \rightarrow$ Particles
- Root Bin Sum \rightarrow Equilibrium Moments
- ∂M_i from $M_i^{Bin} - M_i^{Eq}$
- B2B Collision Recast with δw Particles
- Collision Work $\propto \partial f$, not f
- Valid at Adaptive Tree Depths
- Entropy Estimate for DOF Distribution

Octree ∂f -Boltzmann Bin-to-Bin Collisions



Sample Collisions using
 Δw : for P-P, P-N, N-N, P-M, N-M
M-M: \emptyset




FUTURE DIRECTIONS: HYBRID ∂f

The ∂f -Boltzmann for $f = f^{eq} + f^{dev}$

- ∂f - Concept Old
- How to make ∂f cheaper than full- f (Must adapt DOF Usage..?)
- Recent Progress using $\pm\delta$ -weight Particles
- Requires Remapping due to Particle Growth
- SWPM has Similar Issue (Basis for Octree B2B Collisions)
- Octree: Bin Moments, $M_i \rightarrow$ Particles
- Root Bin Sum \rightarrow Equilibrium Moments
- ∂M_i from $M_i^{Bin} - M_i^{Eq}$
- B2B Collision Recast with δw Particles
- Collision Work $\propto \partial f$, not f
- Valid at Adaptive Tree Depths
- Entropy Estimate for DOF Distribution

Octree ∂f -Boltzmann Bin-to-Bin Collisions



Sample Collisions using
 Δw : for P-P, P-N, N-N, P-M, N-M
 M-M: \emptyset

Combine with 2^6 -Tree
 XV \rightarrow Multigrid Solves?



CONCLUSION

Current Results:

- SWPM+Octree Option for Variable Weight Mixture Collisions
- Multiple Octree Merge only Modification for Multi-Species
- Initial Verification vs. Standard Shock Cases
- Merge/Target Enables Direct Control of Noise
- Unidentified Systematic Error with 1:1 Target
- Initial Testing on 3D Mixture Expansion Better with SWPM+Octree

Future Efforts:

- Additional Investigation of Error Source for Disparate Weights
- Improved Merge/Control of Tails
- Apply to Reacting Flow
- Adaptation for δf



END



Thank You

This material is based upon work supported by the Air Force Office of Scientific Research under award number FA9550-17RQCOR465.

Any opinions, finding, and conclusion or recommendations expressed in this material are those of the author and do not necessarily reflect the views of the United States Air Force.

Questions?